

University of Groningen

Response of *Arabidopsis thaliana* to elevated CO₂ en SO₂

van der Kooij, Thomas Alexander Willem

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2000

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Kooij, T. A. W. (2000). Response of *Arabidopsis thaliana* to elevated CO₂ en SO₂. Groningen: s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ificance for carbon flow in

Van der Kooij T.A.W. & De
Responses of Plants to Air
ys Publishers, Leiden, The

V. 1991. CO₂-Enhanced yield
ated CO₂ environments. Plant

lation of the expression of
regulation by carbohydrates: a
CO₂? Plant Cell Environ. 17:

Summary

Industrial activities and excessive burning of fossil fuel around the world has resulted in a substantial increase of the atmospheric CO₂ and SO₂ concentration. In this thesis the response of *Arabidopsis thaliana* and *Senecio vulgaris* to elevated atmospheric concentrations of CO₂ and SO₂ was investigated. The central issue investigated in this study was the effect of both gaseous air-pollutants on plant metabolism and fitness, with regard to the plants life strategy as a fast growing annual with a short life cycle.

In Chapter 2 *Arabidopsis thaliana* was exposed to an elevated CO₂ concentration of 700 $\mu\text{l l}^{-1}$ CO₂ beginning directly after germination until the end of the life cycle. An increase in dry mass production of 56% was observed in comparison to plants growing at ambient CO₂ concentration of 350 $\mu\text{l l}^{-1}$. The stimulation of the structural biomass production must be ascribed to a short transient stimulation of the relative growth rate (RGR), lasting for one to maximally two days. Thereafter, a similar RGR was observed at both CO₂ concentrations. The rate of photosynthesis was stimulated at elevated CO₂ and no acclimation of photosynthesis was observed after prolonged exposure. Dry matter content was increased at elevated CO₂ which could solely be attributed to the increase of the leaf starch content. Nitrogen content was lower in plants growing at elevated CO₂ but this was solely due to the increase in leaf starch content and the nitrogen content expressed on a fresh weight basis was hardly affected. The combined stimulation of structural biomass and starch content resulted in a substantial increase of the plant fitness reflected in a 51% increase in seed production.

In Chapter 3 *A. thaliana* was exposed to various concentrations of CO₂, ranging from 390 to 1680 $\mu\text{l l}^{-1}$. Maximal stimulation of the shoot structural biomass production was already reached at a CO₂ concentration of 560 $\mu\text{l l}^{-1}$. However, the accumulation of starch continued until the highest applied concentration of 1680 $\mu\text{l l}^{-1}$. The content of soluble sugars and soluble phenolics were not affected by elevated CO₂, except for the highest applied concentration. Photosynthetic capacity was not affected by elevated CO₂ exposure and also the chlorophyll content was hardly affected. The storage capacity of assimilatory starch in the shoot of *A. thaliana* is high and this may be involved in the absence of any disturbance of carbohydrate metabolism and photosynthesis at elevated CO₂.

Thirteen different lines of *A. thaliana* from a world-wide origin were tested for their response to exposure to an elevated CO₂ concentration of 700 $\mu\text{l l}^{-1}$ in Chapter 4. All *A. thaliana* lines show an increase in biomass production, an increase in starch content and a hardly affected soluble sugar content. The latter seems to be a species characteristic. Intra-specific variation existed in the absolute amount of biomass stimulation at elevated CO₂. Part of these differences might be explained by a combination of different factors since lines with thicker leaves

seem to profit less from elevated CO₂ and slower growing lines tend to profit more from the elevated CO₂ than faster growing lines did.

In Chapter 5 the SO₂ response of *A. thaliana* to exposure to SO₂, which is a potent phytotoxic gas, was investigated. *A. thaliana* appeared to be rather SO₂ tolerant. Over a large range of SO₂ concentration up to 0.7 µl l⁻¹ no negative effects on plant growth were observed. A linear relation was observed between the rate of deposition and the atmospheric SO₂ concentration, which resulted in a substantial accumulation of sulfur in the shoot. The main part, 75%, of this absorbed atmospheric sulfur was accumulated as sulfate. The other part could be revealed in the organic sulfur fraction. This 3:1 distribution in sulfate to organic sulfur was irrespective of the applied concentration. The sulfur to nitrogen ratio of the organic fraction was hardly affected indicating no drastic changes in composition or specific accumulation of sulfur rich compounds. Although the shoot content of water-soluble non-protein thiols and glucosinolates was higher upon exposure to SO₂ their contribution to the detoxification of SO₂ was only small. The linear relation between the uptake of SO₂ and the applied concentration was irrespective of the air temperature and only depending on the stomatal conductance as presented in Chapter 6. In contrast the uptake of H₂S showed saturation kinetics.

In Chapter 7 the potential rate of sulfur metabolism was varied by varying the nitrogen availability of *S. vulgaris*, which is a species more susceptible for SO₂ than *A. thaliana*. The uptake rate of SO₂ by *S. vulgaris* were comparable to *A. thaliana* but *S. vulgaris* accumulated to a large extent thiols. Plant biomass production was negatively affected at concentrations above 0.2 µl l⁻¹ SO₂. A substantial increase of sulfate content was observed almost linear to the applied SO₂ concentration and irrespective of the nitrogen nutrition level. The relative decrease in plant biomass production as a result of the SO₂ exposure was irrespective of the applied nitrogen nutrition level. Apparently, the potential to incorporate excessive atmospheric deposited sulfur into organic sulfur at the higher nitrogen nutrition levels did not affect the phytotoxic response towards SO₂ exposure of *S. vulgaris*.

In Chapter 8 thirteen different lines of *A. thaliana* with a world-wide origin were tested in their response to 0.7 µl l⁻¹ SO₂ in order to get more insight into the variability in response to SO₂ within the species. In general the responses of the different lines were similar except for one line, N922, originating from Tadjikistan, which showed an acute phytotoxic response negatively affecting growth and inducing leaf necrosis. All lines accumulated sulfate to a large extent and had a relatively small disturbance of the thiol content. SO₂ tolerance seems to be a species characteristic for *A. thaliana*.

As a ruderal plant species, normally exposed to limiting vegetative expansion and continuous resource capture, *A. thaliana* exhibits an enormous capacity to accumulate available resources during its short vegetative life span. This might

explain the high vacuole capacity, respectively, without a significant increase in addition, *A. thaliana* seems to have a high SO₂ tolerance. The positive response to sulfur composition.

Summary

ing lines tend to profit
ure to SO_2 , which is a
eared to be rather SO_2
0.7 $\mu\text{l l}^{-1}$ no negative
was observed between
ion, which resulted in a
ain part, 75%, of this
The other part could be
on in sulfate to organic
ulfur to nitrogen ratio of
no drastic changes in
mpounds. Although the
ucosinolates was higher
cation of SO_2 was only
he applied concentration
ending on the stomatal
uptake of H_2S showed

as varied by varying the
ore susceptible for SO_2
were comparable to *A.*
t thiols. Plant biomass
bove 0.2 $\mu\text{l l}^{-1}$ SO_2 . A
ost linear to the applied
ition level. The relative
the SO_2 exposure was
arently, the potential to
o organic sulfur at the
ic response towards SO_2

world-wide origin were
et more insight into the
eral the responses of the
1922, originating from
ase negatively affecting
sulfate to a large extent
 SO_2 tolerance seems to

ng vegetative expansion
n enormous capacity to
ve life span. This might

explain the high vacuolar and plastid accumulation of sulfate and starch, respectively, without any negative effects on plant growth and functioning. In addition, *A. thaliana* seems to be a stress tolerator as might be concluded from its high SO_2 tolerance. Therefore, its life strategy might explain its indifferent or even positive response to an anthropogenic induced change of the atmospheric composition.